

## New 1-systems of $Q(6, q)$ , $q$ even

Deirdre Luyckx<sup>1</sup>

Joint work with J. A. Thas

Ghent University, Dept. of Pure Mathematics and Computer Algebra  
Galglaan 2, B-9000 Ghent, Belgium  
dluyckx@cage.rug.ac.be

A 1-system  $\mathcal{M}$  of the parabolic quadric  $Q(6, q)$  in  $\text{PG}(6, q)$  is a set  $\{L_0, L_1, \dots, L_{q^3}\}$  consisting of  $q^3 + 1$  lines on  $Q(6, q)$  having the property that the tangent space of  $Q(6, q)$  at  $L_i$  has no point in common with  $(L_0 \cup L_1 \cup \dots \cup L_{q^3}) \setminus L_i$ ,  $i = 0, 1, \dots, q^3$ . We will discuss a method to construct new locally hermitian 1-systems of  $Q(6, q)$ ,  $q$  even; for  $q$  odd, this was already done in [1]. One of these 1-systems is the spread of the hexagon  $H(q)$ ,  $q = 2^{2e}$ , which was discovered by A. Offer in [3]. Moreover, we can classify these new 1-systems as the only ones on  $Q(6, q)$  which are locally hermitian and semiclassical, but not contained in a 5-dimensional subspace.

Our class of new 1-systems has beautiful applications in a wide range of fields. By projection from the nucleus of  $Q(6, q)$  onto a  $\text{PG}(5, q)$  not containing the nucleus, every 1-system of  $Q(6, q)$ ,  $q$  even, yields a 1-system of  $W_5(q)$ , hence we have also found a new class of 1-systems of  $W_5(q)$ . In [2], it is explained that every 1-system of  $W_5(q)$  yields a semipartial geometry, while by a corollary in [4], a 1-system of  $W_5(q)$  defines a strongly regular graph and a two-weight code. So our new class of 1-systems provides us with new examples of semipartial geometries, strongly regular graphs and two-weight codes.

## References

- [1] D. Luyckx and J. A. Thas, Flocks and locally hermitian 1-systems of  $Q(6, q)$ , In *Finite Geometries - Proceedings of the Fourth Isle of Thorns Conference*, volume 3 of *Developments in Mathematics*, 2001, 257–275.
- [2] D. Luyckx,  $m$ -systems of polar spaces and SPG reguli, *Bulletin of the Belgian Mathematical Society – Simon Stevin*, 2001, to appear.
- [3] A. Offer, Spreads and Ovoids of the Split Cayley Hexagon, University of Adelaide, 2000, second edition, available at <http://cage.rug.ac.be/~nick/Theses/theses.html>
- [4] E.E. Shult and J.A. Thas,  $m$ -systems of polar spaces, *J. Combin. Theory Ser. A* **68** (1994), 184–204.

---

<sup>1</sup>The author is Research Assistant of the Fund for Scientific Research – Flanders (Belgium) (F.W.O.)